

Double parton distributions of the pion

Wojciech Broniowski

17 April 2020

WB, **Enrique Ruiz Arriola**, PRD 101 (2020) 014019 (arXiv:1910.03707)

2018/31/B/ST2/01022



NATIONAL SCIENCE CENTRE
POLAND

- Recent status of double parton scattering (DPS)
- Theoretical issues of double parton distributions (dPDF)
- Simple example of the pion
- Lattice QCD prospects


Motivation for multi-parton distributions

- Old story ([CERN AFS 1986](#), [UA2](#), [Fermilab](#)), renewed interest since ATLAS measurement for $pp \rightarrow W+2 \text{ jets}$ 2013 [Kuti, Weisskopf 1971, Konishi, Ukwu, Veneziano 1979, Gaunt, Stirling 2010, Diehl, Ostermeier, Schäfer 2012, . . . , reviews: Bartalani et al. 2011, Snigirev 2011, Rinaldi, Ceccopieri 2018]
- Model exploration [MIT bag: Chang, Manohar, Waalewijn 2013], valon [WB, ERA, Golec-Biernat 2013, 2016], constituent quarks: Rinaldi, Scopetta, Traini, Vento 2013, 2018]
- Questions: factorization, interference of single- and double-parton scattering [Manohar, Waalewijn 2012, Gaunt 2013], longitudinal-transverse decoupling, positivity bounds [Diehl, Casemets 2013], transverse momentum dependence [Casemets, Gaunt 2019]
- Gaunt-Stirling sum rules [Gaunt, Stirling 2010, WB, ERA 2013, Diehl, Gaunt, Lang, Plößl, Schäfer 2019, 2020]
- Transverse distribution of partons, multi-parton correlations in the hadronic wave function, [background to new physics searches](#) from DPS


Double parton scattering

Example of double parton scattering (DPS)

CMS 1909.06265: **Evidence for WW production from double-parton interactions in pp collisions at $\sqrt{s_{NN}}=13$ TeV**



SPS





DPS

- two additional interactions in SPS $\rightarrow$ cross section suppressed
- absence of jets in DPS $\rightarrow$ handle to reduce the SPS background
- leptonic decays $\rightarrow$ clean final state in the detector

Other DPS searches

CMS: $\sigma_{\text{DPS}, WW} = 1.41 \pm 0.28(\text{stat}) \pm 0.28(\text{syst}) \text{ pb}$ (3.9 σ level)
W+2jets, 2J/ ψ , 2W, 4jets, 3jets+ γ , open charm, ...
[example of predictions from Łuszczak, Maciuła, Szczurek 2011]



DPS can be comparable to SPS at the LHC

Master formula for DPS production

$$\sigma_{hh' \rightarrow AB}^{DPS} = \frac{m}{2} \sum_{i,j,k,l} \int dx_1 dx_2 dx'_1 dx'_2 d^2 b_1 d^2 b_2 d^2 b \times$$

$$D_h^{ij}(x_1, x_2, \mathbf{b}_1, \mathbf{b}_2; \mu) D_{h'}^{kl}(x'_1, x'_2, \mathbf{b} + \mathbf{b}_1, \mathbf{b} + \mathbf{b}_2; \mu) \sigma_A^{ik}(x_1, x'_1; \mu) \sigma_B^{jl}(x_2, x'_2; \mu)$$

($m = 1$ for $A = B$ or 2 otherwise)

If factorization holds ($D_h^{ij}(x_1, x_2, \mathbf{b}_1, \mathbf{b}_2) = D_h^i(x_1) D_h^j(x_2) f_h^i(\mathbf{b}_1) f_h^j(\mathbf{b}_2)$):

$$\sigma_{hh' \rightarrow AB}^{DPS} = \frac{m}{2} \sum_{i,j,k,l} \int d\mathbf{x}_1 d\mathbf{x}_2 d\mathbf{x}'_1 d\mathbf{x}'_2 d^2 \mathbf{q}_\perp \times$$

$$D_h^i(x_1) D_h^j(x_2) F_h^{ij}(\mathbf{q}_\perp) D_{h'}^k(x'_1) D_{h'}^l(x'_2) F_{h'}^{kl}(-\mathbf{q}_\perp) \sigma_A^{ik}(x_1, x'_1) \sigma_B^{jl}(x_2, x'_2)$$

“Pocket” formula

$$\sigma_{hh' \rightarrow AB}^{DPS} \simeq \frac{m}{2} \sigma_{hh' \rightarrow A}^{SPS} \sigma_{hh' \rightarrow B}^{SPS} / \sigma_{\text{eff}}$$

$$\sigma_{\text{eff}} \equiv 1 / \int d^2 q_\perp F_h(\mathbf{q}_\perp) F_{h'}(-\mathbf{q}_\perp)$$

“effective cross section”, typically of the order of the geometric size $\sim 10 - 20$ mb

Double parton distributions

Definition

Intuitive probabilistic definition:

Multi-parton distribution = probability distribution that struck partons have LC momentum fractions x_i

Field-theoretic definition of (spin-averaged) sPDF and dPDF [Diehl, Ostermeier, Schaeffer 2012] of a hadron with momentum p :

$$D_j(x) = \int \frac{dz^-}{2\pi} e^{ixz^-p^+} \langle p | \mathcal{O}_j(0, z) | p \rangle \Big|_{z^+=0, \mathbf{z}=0}$$
$$D_{j_1 j_2}(x_1, x_2, \mathbf{b}) = 2p^+ \int dy^- \frac{dz_1^-}{2\pi} \frac{dz_2^-}{2\pi} e^{i(x_1 z_1^- + x_2 z_2^-)p^+}$$
$$\times \langle p | \mathcal{O}_{j_1}(y, z_1) \mathcal{O}_{j_2}(0, z_2) | p \rangle \Big|_{z_1^+ = z_2^+ = y^+ = 0, \mathbf{z}_1 = \mathbf{z}_2 = 0}$$

$$\mathcal{O}_q(y, z) = \frac{1}{2} \bar{q}(y - \frac{z}{2}) \gamma^+ q(y + \frac{z}{2}), \dots \quad (\text{LC gauge}) \quad v^\pm = (v^0 \pm v^3)/\sqrt{2}$$


$y = (y^+, y^-, \mathbf{b})$, ($\mathbf{b}$ is the transverse distance between the two quarks)

(no problem with gauge links!)

dPDF in momentum space

Fourier transform in $\mathbf{b}$

$$D_{j_1 j_2}(x_1, x_2, \mathbf{b}) \rightarrow \tilde{D}_{j_1 j_2}(x_1, x_2, \mathbf{q})$$



Special case of $\mathbf{q} = \mathbf{0}$:

$$D_{j_1 j_2}(x_1, x_2) \equiv \tilde{D}_{j_1 j_2}(x_1, x_2, \mathbf{q} = \mathbf{0}) = \int d^2 b D_{j_1 j_2}(x_1, x_2, \mathbf{b})$$

Gaunt-Stirling sum rules

[Gaunt, Stirling, JHEP (2010) 1003:005, Gaunt, PhD Thesis]

Fock-space decomposition on the light front, mom. conservation $\rightarrow$

$$\int_0^{1-x_2} dx_1 D_{i_{\text{val}}j}(x_1, x_2) = (N_{i_{\text{val}}} - \delta_{ij} + \delta_{\bar{i}j}) D_j(x_2) \quad (\text{quark number})$$

$$\sum_i \int_0^{1-x_2} dx_1 x_1 D_{ij}(x_1, x_2) = (1 - x_2) D_j(x_2) \quad (\text{momentum})$$

$$(A_{i_{\text{val}}} \equiv A_i - A_{\bar{i}}) \qquad N_{i_{\text{val}}} = \int_0^1 dx D_{i_{\text{val}}}(x)$$

- Preserved by dDGLAP evolution
- Non-trivial to satisfy with the (guessed) function
- Checked in light-front perturbation theory and in lowest-order covariant calculations in [Diehl, Plöchl, Schäfer 2019]


Important and fundamental constraints!

The pion

Why the pion

- Much simpler theoretically than the proton
- Possible future results for dPDFs from lattice correlators in the pion state, extending the studies of [Zimmermann et al. 2017, 2018] → moments in x_1 and x_2 and the transverse form factor
- Nicely illustrates formalism, theoretical features

Chiral quark models




- χ SB breaking $\rightarrow$ massive quarks
- Point-like interactions
- Soft matrix elements with pions (and photons, W , Z)
- Large- $N_c \rightarrow$ one-quark loop
- Regularization

pion – Goldstone boson of χ SB, fully relativistic $q\bar{q}$ bound state of the Bethe-Salpeter equation

Quantities evaluated at the **quark model** scale
(where **constituent quarks** are the only degrees of freedom)

Chiral quark models



- χ SB breaking $\rightarrow$ massive quarks
- Point-like interactions
- Soft matrix elements with pions (and photons, W , Z)
- Large- $N_c \rightarrow$ one-quark loop
- Regularization


pion – Goldstone boson of χ SB, fully relativistic $q\bar{q}$ bound state of the Bethe-Salpeter equation

Need for QCD evolution, hard-soft factorization

Gluon dressing, renorm-group improved

Creates bulk of the effect

[Davidson, Arriola, 1995]


In the chiral limit of $m_{\pi} = 0$:

$$q_{\text{val}}(x; Q_0) = 1 \times \theta[x(1-x)]$$

Quarks are the only degrees of freedom, hence saturate the PDF sum rules:

$$\int_0^1 dx q_{\text{val}}(x; Q_0) = 1 \text{ (valence)}, \quad 2 \int_0^1 dx x q_{\text{val}}(x; Q_0) = 1 \text{ (momentum)}$$

Pion valence quark sPDF, NJL vs E615



points: Fermilab E615

Drell-Yan, $\pi^\pm W \rightarrow \mu^\pm \mu^\mp X$

dashed line: 2005 NLO
reanalysis [Wijesoorija et al.]

band: QM + LO DGLAP
from $Q_0 = 313_{-10}^{+20}$ MeV to
 $Q = 4$ GeV


Other results: GPD [WB, ERA, Golec-Biernat 2008], TDA [WB, ERA 2007], equal-time wave functions [WB, ERA, Prelovsek, Šantelj 2009], transversity distributions [WB, ERA, Dorokhov 2009], quasi-distribution amplitude [WB, ERA, 2018]

dPDF of the pion in NJL (new stuff)

- WB talk at Light Cone 2019, 16-20 Sep. 2019, Palaiseau, France
- A. Courtoy, S. Noguera, and S. Scopetta, JHEP 12, 045 (2019), arXiv:1909.09530
- WB, ERA, PRD 101 (2020) 014019, arXiv:1910.03707

dPDF of the pion in NJL model

In LC kinematics only one diagram:



In the chiral limit of $m_\pi^2 = 0$ a very simple result follows:

$$D_{u\bar{d}}(x_1, x_2, \mathbf{q}_\perp) = \delta(1 - x_1 - x_2) \theta[x_1(1 - x_1)] \theta[x_2(1 - x_2)] F(\mathbf{q}_\perp^2)$$

momentum conservation


support

form factor

- Factorization (in the chiral limit) of the longitudinal and transverse dynamics
- GS sum rules satisfied (preserved by the evolution)
- Results at the quark-model scale $\rightarrow$ evolution

dDGLAP evolution in the Mellin space

[Kirschner 1979, Shelest, Snigirev, Zinovev 1982]: method of solving dDGLAP based on the Mellin moments, similarly to sPDF **simplification for valence distributions**







$$t = \frac{1}{2\pi\beta} \log [1 + \alpha_s(\mu)\beta \log(\Lambda_{\text{QCD}}/\mu)], \quad \beta = \frac{11N_c - 2N_f}{12\pi}$$

Valence:


$$\text{dPDF : } \frac{d}{dt} M_{j_1 j_2}^{n_1 n_2}(t) = (P_{j_1 \rightarrow j_1}^{n_1} + P_{j_2 \rightarrow j_2}^{n_2}) M_{j_1, j_2}^{n_1 n_2}(t)$$

$$\text{sPDF : } \frac{d}{dt} M_j^n(t) = P_{j \rightarrow j}^n M_j^n(t)$$


$$x_1 x_2 D_{u\bar{d}}^{\pi^+}(x_1, x_2)$$



$D_{u\bar{d}}^{\pi^+}(x_1, x_2) - \log \text{ scale}$



Correlation (cf. [Golec-Biernat et al. 2015] for gluons in p)




Valence moments in NJL

$$\frac{\langle x_1^n x_2^m \rangle}{\langle x_1^n \rangle \langle x_2^m \rangle} = \frac{(1+n)!(1+m)!}{(1+n+m)!} \quad (\text{NJL, any scale})$$

(independent of the evolution scale)

	1	2	3	4
1	2 3	$\frac{1}{3}$	$\frac{2}{5}$	$\frac{1}{3}$
2	$\frac{1}{2}$	$\frac{3}{10}$	$\frac{1}{5}$	$\frac{1}{7}$
3	$\frac{2}{5}$	$\frac{1}{5}$	$\frac{4}{35}$	$\frac{1}{14}$
4	$\frac{1}{3}$	$\frac{1}{7}$	$\frac{1}{14}$	$\frac{5}{126}$



Double moments reduced compared to products of single moments
(longitudinal momentum conservation, **would be 1 if no correlations!**)

Pion on the lattice

sPDF:


Lowest moment of sPDF evaluated on the lattice [Abdel-Rehim et al. 2016]:

$$\langle x \rangle_{\text{val}} = 0.214_{-9}^{+12} \text{ at } \mu = 2 \text{ GeV}, m_\pi = 139 \text{ MeV}$$

$$\langle x \rangle_{\text{val}} = 0.256(13) \text{ phenomenologically [Wijesooriya et al. 2005]}$$

dPDF: not yet, but similar correlations studied [Zimmermann et al. 2017, 2018]

C1



More lattice results expected ...

$$\langle x_1 x_2 \rangle_{\text{val}} / \langle x \rangle_{\text{val}}^2$$

simplest (and to LO scale independent) measure of longitudinal correlations

Transverse structure

- Depends on the regulator, good for low $\mathbf{q}_\perp$
- Transverse form factor in NJL with spectral regularisation (vector meson dominance)

$$F(\mathbf{q}_\perp) = \frac{m_\rho^4 - \mathbf{q}_\perp^2 m_\rho^2}{(m_\rho^2 + \mathbf{q}_\perp^2)^2},$$

- Recall $\sigma_{\text{DPS}}^{AB} = \frac{m}{2} \sigma_{\text{SPS}}^A \sigma_{\text{SPS}}^B / \sigma_{\text{eff}}$
- Effective cross section (here coincides with geometric)

$$\sigma_{\text{eff}} = \frac{1}{\int \frac{d^2 \mathbf{q}_\perp}{(2\pi)^2} F(\mathbf{q}_\perp) F(-\mathbf{q}_\perp)} = \pi \frac{12}{m_\rho^2} = \pi \langle b^2 \rangle_f = 23 \text{ mb}$$

Summary

- Topic driven by recent experimental evidence as well as future lattice studies
- Important issues of correlations breaking the typically assumed factorizations
- NJL: simplest field theory of the pion in the **soft** regime; spontaneous chiral symmetry breaking; covariant calculation, all symmetries preserved → good features
- dPDF in NJL = $\delta(1 - x_1 - x_2) \times F(q_\perp^2) + \text{dDGLAP evolution}$; longitudinal-transverse factorization (in the chiral limit)
- Correlations decrease with increasing evolution scale and are probably not very important ($\pm 25\%$) in the range probed by experiments (which at the moment are concerned with orders of magnitude), justifying the **product ansatz** in that limit
- Ratios of Mellin moments measure the $x_1 - x_2$ **factorization breaking**; could be verified in future lattice calculations
- The effective cross section in the model is **geometric**

Back-up

