Discontinuous phase transitions in the q-voter model with generalized anticonformity on random graphs. Angelika Abramiuk, Arkadiusz Lipiecki, Jakub Pawłowski and Katarzyna Sznajd-Weron

Department of Theoretical Physics, Faculty of Fundamental Problems of Technology, Wrocław University of Science and Technology

Motivation

- In 2019, a generalization of q-voter model with anticonformity was proposed [1]. The novelty was decoupling of sizes of groups of influence needed for conformity and anticonformity. In result, a previously unseen discontinuous phase transition was observed.
- Discontinuous phase transitions are connected with interesting social phenomena such as tipping points and hysteresis.
- The q-voter model with generalized anticonformity has not been examined on random graphs, which are more realistic from a sociological point of view. Hence, one may ask the question, **does the discontinuous** phase transition survive the transition to random graphs?

Model description

We consider a collection of N voters located in the vertices of a random graph. Each of them is characterized by a binary variable $S_i(t) = \pm 1$, i = $1, \ldots N$, where t denotes time. They interact only with their neighbors, i.e., vertices with a direct link to them.

Dynamics

Model parameters:

- p probability of anticonformity,
- q_a necessary group size for anticonformity,
- q_c necessary group size for anticonformity.
- k average node degree of the network



Pair approximation

In our work, we consider the network topology different from the complete graph, hence the usually employed mean field approximation is insufficient. We need to take into account the correlations between the nearest neighbors. Here, the model is described by two variables, the concentration of positive opinion $c(t) \equiv \frac{N_{\uparrow}(t)}{N}$, and the concentration of active bonds $b(t) = \frac{1}{2Nk} \sum_{i=1}^{N} \sum_{j=1}^{N} (1 - S_i(t)S_j(t)) \mathbf{1}_{(i,j)\in E}$ i.e. bonds between opposite spins. Such a description is possible because of two assumptions:

- the states of agent's neighbors are independent of each other,
- there is no correlation between vertex degree and finding an agent with a given opinion or a bond in a given state

This constitutes the Pair approximation [2], resulting in two coupled differential equations to be solved by numerical methods - see [3].







Conclusion

• We implemented the q-voter model with generalized anticonformity on random E-R graphs and conducted Monte Carlo simulations.

• It occurred that the theoretical predictions obtained via Pair Approximation do not match the simulation results for values of q_a fulfilling: $q_a \ge q_c - 1.$

• In these cases, PA yields a discontinuous phase transition and nonzero width of the hysteresis, whereas Monte Carlo simulations produce a continuous phase transition, the same as MFA. • To conclude, PA should be used with caution, and to ensure the validity of the results, it is best to complement it with simulation results.

References

[1] A Abramiuk, J Pawłowski, and K Sznajd-Weron. Is independence necessary for a discontinuous phase transition within the q-voter model? Entropy, 21(5), 2019.

[2] A Jędrzejewski.

Pair approximation for the q -voter model with independence on complex networks. *Physical Review E*, 95(1):1–9, 2017.

[3] A Abramiuk, A Lipiecki, J Pawłowski, and K Sznajd-Weron.

Discontinuous phase transitions in the q-voter model with generalized anticonformity on random graphs.

Manuscript submitted for publication., 2021.

Acknowledgements

This work has been partially supported by the National Science Center (NCN, Poland) through grants no. 2016/21/B/HS6/01256 and 2019/35/B/HS6/02530, and by Polish Ministry of Science and Higher Education through project "Diamentowy Grant" no. DI2019 0150 49. More extensive computations have been conducted using the PLGrid Infrastructure.



Wrocław University of Science and Technology